

Light Curves and Spectra of Dust Echoes From Gamma-Ray Bursts and their Afterglows: Continued Evidence that GRB 970228 is Associated with a Supernova

Daniel E. Reichart^{1,2}

ABSTRACT

The relative brightening and reddening of the optical afterglows of GRB 970228 and GRB 980326 around 20 – 30 days after these bursts have been attributed to supernovae, which are red (blueward of their spectral peak), and which peak in time after $\approx 20(1+z)$ days. However, this direct evidence for a GRB/SN connection has recently been challenged. It has been suggested that the late afterglows of these bursts can be explained by dust echoes, of which we consider two cases: (1) the scattering of light from the afterglow (the forward shock), the optical flash (the reverse shock), and/or optical light from the burst itself by a shell of dust at a radius R from the progenitor, and (2) the absorption and thermal re-emission of this light by this shell of dust. In this paper, we model and compute dust echo light curves and spectra for both cases. Although the late-time afterglow of GRB 980326 was not sufficiently well sampled to rule out a dust echo description, we find that the late-time afterglow of GRB 970228 cannot be explained by a dust echo.

Subject headings: dust, extinction — gamma-rays: bursts

1. Introduction

Bloom et al. (1999) first identified a second component to the optical afterglow of GRB 980326. This second component grew brighter than the extrapolation of the early afterglow (which faded as a power law) around 1 week after the burst, and peaked around 20 – 30 days after the burst before fading away. Rough spectra taken 2.4 and 28 days after the burst show that the second component was significantly redder than the early afterglow. Bloom et al. (1999) attributed this behavior to a supernova at a redshift of $z \sim 1$. Supernovae peak after about

¹Department of Astronomy, California Institute of Technology, Mail Code 105-24, 1201 East California Boulevard, Pasadena, CA 91125

²Hubble Fellow

$20(1+z)$ days and are significantly redder than afterglows (blueward of the supernova’s spectral peak). Unfortunately, neither a redshift nor multiband photometry is available for this afterglow.

Reichart (1999) identified a similar second component to the optical afterglow of GRB 970228, the first burst for which an afterglow was detected. Reichart (1999) and Galama et al. (2000) showed that the light curve of this second component is consistent with the expectation of a supernova at the spectroscopically-measured redshift of this burst ($z = 0.695$; Bloom, Djorgovski, & Kulkarni 2001). More convincingly, VRIJK photometry of the second component taken near its peak showed that its spectral flux distribution is also consistent with the expectation of a supernova at this redshift: (1) the spectral flux distribution is very red blueward of the spectral peak and very blue redward of the spectral peak, and (2) the spectral peak is at the frequency of the spectral peak of a supernova at $z \approx 0.7$ (Figure 1; Reichart 1999; Galama et al. 2000; Reichart, Castander, & Lamb 2000; Reichart, Lamb, & Castander 2000). This is the best evidence to date for the GRB/SN connection.

The GRB/SN connection has recently been challenged by alternative explanations of these observations, called dust echoes (Waxman & Draine 2000; Esin & Blandford 2000; see also Lamb 1997, 2000). If the progenitors of bursts are massive stars (see Lamb & Reichart 2000 for a review of the indirect evidence that bursts are associated with star-forming regions), due to the progenitor’s wind, the density of the surrounding medium should fall off as $n(r) \propto r^{-2}$ out to the radius R_w of the wind termination shock (Chevalier & Li 1999). At R_w , the density should jump by a factor of ≈ 4 , and beyond R_w , the density should remain approximately constant out to great distances (Chevalier & Li 1999). Dust in the vicinity of the burst should be sublimated by UV light from the optical flash (the reverse shock; Waxman & Draine 2000). Let R be the radius out to which dust is sublimated. If $R < R_w$, and if the dust is sublimated isotropically, the dust should occupy a relatively thin shell at $\approx R$ due to the rapidly decreasing density of the circumburst medium with distance. Light that is emitted by the afterglow (the forward shock) from a distance that is much less than R at an angle θ relative to the line of sight, is scattered (Esin & Blandford 2000) or absorbed and thermally re-emitted (Waxman & Draine 2000; Esin & Blandford 2000) by dust at $r \approx R$ into the line of sight, and arrives at the observer, arrives with an observer-frame time delay of

$$t = \frac{r(1+z)}{c}(1 - \cos \theta) \quad (1)$$

relative to light that was emitted into the line of sight ($\theta = 0$) and not scattered or absorbed. This dust and light geometry is depicted in the top panel of Figure 2. Waxman & Draine (2000) and Esin & Blandford (2000) find that this results in secondary components to the afterglow, called dust echoes, that may peak about 20 – 30 days after the burst, and that may be significantly redder than the early afterglow, i.e., that the late-time afterglows of GRB 970228 and GRB 980326 may in fact be due to dust echoes, and not supernovae.

There are a number of problems with this dust and light geometry. First of all, $R \approx 12$ pc (Waxman & Draine 2000); however, R_w is probably only a few parsecs (Chavalier & Li 2000).

Consequently, the dust would probably not occupy a thin shell.

Secondly and more importantly, UV light from the optical flash will only sublimate dust within θ_{jet} of the line of sight, where θ_{jet} is the initial half-opening angle of the jet. (The Lorentz factor γ of the shocked material in the jet is much greater than θ_{jet}^{-1} when the optical flash occurs.) If the shock is adiabatic,

$$\gamma \approx 6.1(1+z)^{3/8} \left(\frac{n}{1 \text{ cm}^3} \right)^{-1/8} \left(\frac{E}{10^{52} \text{ erg}} \right)^{1/8} \left(\frac{t}{1 \text{ dy}} \right)^{-3/8}, \quad (2)$$

where n is the number density of gas atoms in the circumburst medium, E is the isotropic-equivalent energy of the shock, and t is observer-frame time (e.g., Sari, Piran, & Narayan 1998). Let $n = n(1 \text{ pc})(r/1 \text{ pc})^{-2}$. Since $r \approx 4\gamma^2 ct/(1+z)$ (e.g., Sari, Piran, & Narayan 1998),

$$\gamma \approx 2.2(1+z)^{1/4} \left[\frac{n(1 \text{ pc})}{1 \text{ cm}^3} \right]^{-1/4} \left(\frac{E}{10^{52} \text{ erg}} \right)^{1/4} \left(\frac{t}{1 \text{ dy}} \right)^{-1/4}. \quad (3)$$

The afterglow emits light into a half-opening angle $\theta_{col} \approx (\theta_{jet}^2 + \gamma^{-2})^{1/2}$. Given the rate at which γ decreases with time, θ_{col} quickly grows significantly greater than θ_{jet} : For example, $\theta_{col} > 1.1\theta_{jet}$ after only a few minutes if $\theta_{jet} \approx 0.1$:³

$$t \gtrsim 14(1+z) \left[\frac{n(1 \text{ pc})}{1 \text{ cm}^3} \right]^{-1} \left(\frac{E\theta_{jet}^2}{10^{52} \text{ erg}} \right) \left(\frac{\theta_{jet}}{0.1} \right)^2 \text{ min}. \quad (4)$$

Light emitted at $\theta > \theta_{jet}$ would be scattered, absorbed, and thermally re-emitted by dust at radii that are substantially smaller than R . Due to the substantially higher density of dust at these radii, this light would dominate the dust echo. Radiative shocks yield similar results. This more realistic dust and light geometry is depicted in the bottom panel of Figure 2.

In this paper we consider a dust and light geometry of which the dust and light geometry depicted in the bottom panel of Figure 2 is a limiting case. High-energy light from the progenitor may destroy dust (e.g., via electrostatic disruption; Draine & Salpeter 1979; Waxman & Draine 2000) out to a smaller radius. Similarly, irregularities in the progenitor’s wind may destroy dust in shocks. In either case, the dust would be destroyed isotropically. Let R instead be the radius out to which dust is destroyed by the progenitor, and let R_{jet} be the radius out to which dust is destroyed by the burst, the optical flash, and/or the afterglow, by whatever dust destruction mechanisms. If R_{jet} is even twice as great as R , and if the jet is oriented along the line of sight, the dust echoes should brighten substantially at a time given by Equation (1) with $r \approx R$ and

³When $\gamma \approx \theta_{jet}^{-1}$, the jet begins to expand laterally, and the afterglow fades much more rapidly (e.g., Panaitescu, Mészáros, & Rees 1998). Applying Equation (3) to the afterglows of GRB 990510 (e.g., Stanek et al. 1999; Harrison et al. 1999), GRB 991216 (Halpern et al. 2000), GRB 000301c (e.g., Rhoads & Fruchter 2000; Berger et al. 2000), and GRB 000926 (Price et al. 2001), we find that $\theta_{jet} \approx 0.17[n(1 \text{ pc})/1 \text{ cm}^3]^{1/4}$, $0.14[n(1 \text{ pc})/1 \text{ cm}^3]^{1/4}$, $0.38[n(1 \text{ pc})/1 \text{ cm}^3]^{1/4}$, and $0.18[n(1 \text{ pc})/1 \text{ cm}^3]^{1/4}$, respectively.

$\theta \approx \theta_{jet}$, and dim substantially at a time given by Equation (1) with $r \approx R$ and $\theta \approx \pi - \theta_{jet}$, since we assume that the jet is bipolar. If $\theta_{jet} \ll 1$, then the dust echoes “turn on” at

$$t_{on} \approx 6.0(1+z) \left(\frac{R}{1 \text{ pc}} \right) \left(\frac{\theta_{jet}}{0.1} \right)^2 \text{ dy}, \quad (5)$$

and “turn off” at

$$t_{off} \approx 6.5(1+z) \left(\frac{R}{1 \text{ pc}} \right) \text{ yr} - t_{on}, \quad (6)$$

though they may fade earlier than t_{off} due to other effects (see §2 and §3). The jet may be oriented as much as $\approx \theta_{jet}$ from the line of sight (any more and we would not have seen the burst). Consequently, the dust echoes may “turn on” half of the way as early as $t \approx 0$ and all of the way as late as $t \approx 4t_{on}$. The “turn off” time would be similarly affected. The dust and light geometry depicted in the bottom panel of Figure 2 is recovered in the limit that $R \rightarrow 0$.

Another dust and light geometry that we consider in this paper is that light from the optical flash, as well as optical light that may be associated with the burst itself, may result in dust echoes. Consequently, dust echoes may be a way to measure the brightness of the optical flash and/or the optical brightness of the burst.

In this paper, we model and compute dust echo light curves and spectra for both $R_{jet} > a$ few times R , and $R_{jet} < R$ (in which case dust is destroyed only isotropically). Any other case would be fine tuning. We also model and compute dust echoes resulting from a pulse of light (i.e., the optical flash or optical light associated with the burst), and an afterglow light curve. Finally, we model and compute dust echo light curves and spectra for both scattered light (§2), and absorbed and thermally re-emitted light (§3). We find that both types of dust echoes have light curves and spectra that can be distinguished from each other, and from the light curves and spectra of supernovae. In §4, we re-examine the late-time afterglow of GRB 970228, and find that it cannot be explained by a dust echo. We draw conclusions in §5.

2. Dust Echoes from Scattered Light

2.1. Dust Echoes from a Scattered Isotropic Pulse of Light

Before modeling the general case of the dust echo of collimated light that is emitted as a function of time (e.g., an afterglow), we first model the simpler case of the dust echo of an isotropic pulse of light (e.g., an optical flash or optical light associated with the burst) that is emitted from a distance that is much less than R . Consider the light that is emitted at an angle θ relative to the line of sight, is scattered by dust at r into the line of sight, and arrives at the observer. The optical depth at observer-frame frequency ν along this path is given by:

$$\tau_{\nu(1+z)}(r, \theta) = \int_0^r n(r, \theta) \sigma_{\nu(1+z)}^{tot} dr + \int_{r \cos \theta}^{\infty} n(r, \theta) \sigma_{\nu(1+z)}^{tot} d(r \cos \theta), \quad (7)$$

where $n(r, \theta)$ is the number density of dust particles at r and θ , and $\sigma_{\nu(1+z)}^{tot} = \sigma_{\nu(1+z)}^{abs} + \sigma_{\nu(1+z)}^{sc}$ is the total (absorption plus scattering) cross section of these particles. The number density of dust particles at $r < R_w$ is given by (§1):

$$n(r, \theta) = \begin{cases} 0 & (r < R) \\ 0 & (r < R_{jet} \text{ and } 0 < \theta < \theta_{jet} \text{ or } \pi - \theta_{jet} < \theta < \pi) \\ n_R(r/R)^{-2} & (R < r < R_w \text{ and } f\theta_{jet} < \theta < \pi - f\theta_{jet}) \\ n_R(r/R)^{-2} & (R_{jet} < r < R_w \text{ if } R < R_{jet} < R_w) \end{cases}, \quad (8)$$

where (§1)

$$f = \begin{cases} 0 & (R_{jet} < R) \\ 1 & (R_{jet} > \text{a few times } R) \end{cases}. \quad (9)$$

Assuming that $R_w > \text{a few times } R$, we find that

$$\tau_{\nu(1+z)}(r, \theta) = \begin{cases} \tau_{\nu(1+z)} \left[1 + \frac{R}{r} \left(\frac{\theta - f\theta_{jet}}{\sin \theta} - 1 \right) \right] & (f\theta_{jet} < \theta < \frac{\pi}{2} \text{ or } r \sin \theta > R) \\ \tau_{\nu(1+z)} \left[1 + \frac{R}{r} \left(\frac{\theta - f\theta_{jet} - 2 \cos^{-1} \frac{r \sin \theta}{R}}{\sin \theta} - 1 \right) \right] & (\frac{\pi}{2} < \theta < \pi - f\theta_{jet} \text{ and } r \sin \theta < R) \\ \tau_{\nu(1+z)} (3 - \frac{2R}{r}) & (\theta = \pi \text{ and } f = 0) \end{cases}, \quad (10)$$

where $\tau_{\nu(1+z)} = n_R \sigma_{\nu}^{tot} R$ is the optical depth along any line of sight if $f = 0$, or any line of sight between $\theta_{jet} < \theta < \pi - \theta_{jet}$ if $f = 1$.

The probability that light takes this path, i.e., that light that is emitted within $d\Omega$ of θ makes it to within dr of r without being scattered or absorbed, is scattered within dr of r into the line of sight, and makes it to the observer without being scattered or absorbed, is given by:

$$dP_{\nu(1+z)}^{sc}(r, \theta) = e^{-\tau_{\nu(1+z)}(r, \theta)} n(r, \theta) \frac{d\sigma_{\nu(1+z)}^{sc}}{d\Omega} d\Omega dr. \quad (11)$$

We adopt the same differential dust scattering cross section as do Esin & Blandford (2000; White 1979):

$$\frac{d\sigma_{\nu(1+z)}^{sc}}{d\Omega} = \frac{\epsilon_{\nu(1+z)} \sigma_{\nu(1+z)}^{tot}}{4\pi} \frac{1 - \bar{\mu}_{\nu(1+z)}^2}{[1 + \bar{\mu}_{\nu(1+z)}^2 - 2\bar{\mu}_{\nu(1+z)}\mu]^{3/2}}, \quad (12)$$

where $\mu = \cos \theta$, $\bar{\mu}_{\nu(1+z)}$ is the mean value of μ at $\nu(1+z)$ for the dust particles, and $\epsilon_{\nu(1+z)} = \sigma_{\nu(1+z)}^{sc} / \sigma_{\nu(1+z)}^{tot}$ is the mean albedo at $\nu(1+z)$ for these particles. We adopt $\bar{\mu}_{\nu(1+z)} = \epsilon_{\nu(1+z)} = 0.5$ (White 1979). The solid angle across which this light is scattered is $d\Omega = 2\pi \sin \theta d\theta$. Consequently,

$$dP_{\nu(1+z)}^{sc}(r, \theta) = \frac{3\tau_{\nu(1+z)}}{2} \frac{n(r, \theta)}{n_R} \frac{e^{-\tau_{\nu(1+z)}(r, \theta)} \sin \theta d\theta}{(5 - 4 \cos \theta)^{3/2}} d\left(\frac{r}{R}\right). \quad (13)$$

Using Equation (1), we rewrite Equation (13) as a function of r and t :

$$dP_{\nu(1+z)}^{sc}(r, t) = \frac{3c\tau_{\nu(1+z)}}{2R(1+z)} \frac{n(r, \theta)}{n_R} \left(\frac{r}{R}\right)^{-1} \frac{e^{-\tau_{\nu(1+z)}(r, \theta)}}{(5 - 4 \cos \theta)^{3/2}} d\left(\frac{r}{R}\right) dt, \quad (14)$$

where θ is a function of r and t given by Equation (1).

Let $F_\nu dt$ be the fluence of the pulse of light at ν . Ignoring light that multiply scatters into the line of sight, the light curve of the dust echo is then given by:

$$F_\nu^E(t) = \int \frac{F_\nu}{e^{-(1-f)\tau_\nu(1+z)}} dP_{\nu(1+z)}^{sc}(r, t) \quad (15)$$

$$= \frac{3c\tau_\nu(1+z)}{2R(1+z)} \int_r^\infty P(r') \left(\frac{r'}{R}\right)^{-3} \frac{e^{-[\tau_\nu(1+z)(r', \theta) - (1-f)\tau_\nu(1+z)]}}{(5 - 4 \cos \theta)^{3/2}} d\left(\frac{r'}{R}\right) F_\nu dt, \quad (16)$$

where

$$P(r') = \begin{cases} 0 & [r' < r(\pi - f\theta_{jet})] \\ 1 & [r(\pi - f\theta_{jet}) < r' < r(f\theta_{jet})] \\ 0 & [r' > r(f\theta_{jet})] \end{cases}, \quad (17)$$

$$r = \begin{cases} R & [t < t(\pi - f\theta_{jet})] \\ r(\pi - f\theta_{jet}) & [t > t(\pi - f\theta_{jet})] \end{cases}, \quad (18)$$

and

$$\theta = \begin{cases} \theta(R) & [t < t(\pi - f\theta_{jet})] \\ \pi - f\theta_{jet} & [t > t(\pi - f\theta_{jet})] \end{cases}, \quad (19)$$

and where

$$r(\theta) = \frac{ct}{(1 - \cos \theta)(1 + z)}, \quad (20)$$

$$\theta(r) = \cos^{-1} \left[1 - \frac{ct}{r(1 + z)} \right], \quad (21)$$

and

$$t(\theta) = \frac{(1 - \cos \theta)R(1 + z)}{c}. \quad (22)$$

If $f = 0$, or if $f = 1$ and $\theta_{jet} \ll \min\{\theta, (\theta/\tau)(r/R)\}$ [Equations (17) and (10), respectively], then Equation (16) is approximately given by:

$$F_\nu^E(t) \approx \begin{cases} A \left(\frac{r}{R}\right)^{-2} & (t < t_{sc} \text{ and } t < t_\tau) \\ A \left(\frac{r}{R}\right)^{-2} \left(\frac{t}{t_{sc}}\right)^{-3/2} \left[4 - 3 \left(\frac{t}{t_{sc}}\right)^{-1/2} \right] & (t_{sc} < t < t_\tau) \\ A \left(\frac{r}{R}\right)^{-2} \left(\frac{t_\tau}{t_{sc}}\right)^{-3/2} \left(\frac{t}{t_\tau}\right)^{-7/4} \left[4 - 3 \left(\frac{t_\tau}{t_{sc}}\right)^{-1/2} \left(\frac{t}{t_\tau}\right)^{-1/4} \right] & (t_{sc} < t_\tau < t) \\ A \left(\frac{r}{R}\right)^{-2} \left(\frac{t}{t_\tau}\right)^{-1} & (t_\tau < t \text{ and } t_\tau \ll t_{sc}) \end{cases}, \quad (23)$$

where

$$A = \frac{3c\tau_\nu(1+z)}{4R(1+z)} e^{-f\tau_\nu(1+z)} F_\nu dt, \quad (24)$$

$$t_{sc} = \frac{1}{4} \frac{R(1+z)}{c} \left(\frac{r}{R}\right) \quad (25)$$

is the time when the scattering cross section begins to fall off due to scattering at greater angles, and

$$t_\tau \approx \frac{3}{\tau_{\nu(1+z)}} \frac{R(1+z)}{c} \left(\frac{r}{R} \right)^2 \quad (26)$$

is the time when the optical depth grows greater by 1 due to the greater path length at greater scattering angles. This interaction of the dust and light geometries is depicted in the top panel of Figure 3. We plot example light curves for different values of f , $R(1+z)$, and $\tau_{\nu(1+z)}$ in the top panels of Figures 4 – 7; however, consider the following limiting cases:

f = 0 (Figures 4 and 5, top panels): If $\tau_{\nu(1+z)} \lesssim 12$, then $t_{sc} \lesssim t_\tau$. Consequently, $F_\nu^E(t) \propto t^0$ from $t = 0$ until $t \approx t_{sc}$, after which the light curve slowly rolls over to $F_\nu^E(t) \propto t^{-3/2}$. If $\tau_{\nu(1+z)} \lesssim 1.5$, then this continues until $t \approx t_{off}$, after which $F_\nu^E(t) \propto t^{-2}$. If $1.5 \lesssim \tau_{\nu(1+z)} \lesssim 12$, then this continues until $t \approx t_\tau$, after which the light curve slowly rolls over to $F_\nu^E(t) \propto t^{-7/4}$ until after $t = t_{off}$. After this time, $F_\nu^E(t) \propto t^{-2}$. If $\tau_{\nu(1+z)} \gg 12$, then t_τ is initially less than t_{sc} . Consequently, $F_\nu^E(t) \propto t^0$ from $t \approx 0$ until $t \approx t_\tau$, after which $F_\nu^E(t) \propto t^{-1}$ until after $t = t_{off}$. After this time, $F_\nu^E(t) \propto t^{-2}$.

f = 1 (Figures 6 and 7, top panels): The light curve has the same dependence on time as in the $f = 0$ case except that it does not “turn on” until $\sim t_{on}$. Also, since the dust is destroyed along the line of sight, the pulse of light is a factor of $e^{\tau_{\nu(1+z)}}$ brighter than in the $f = 0$ case. Consequently, the light curve, which is normalized to the fluence of the pulse of light, is a factor of $e^{-\tau_{\nu(1+z)}}$ fainter than in the $f = 0$ case.

Let $\tau_{\nu(1+z)} \propto [\nu(1+z)]^{\alpha_{\nu(1+z)}}$. The value of $\alpha_{\nu(1+z)}$ is canonically assumed to be 1; however, the true value is closer to ≈ 1.6 at NIR values of $\nu(1+z)$, and levels off to ≈ 1 in the UV (Reichart 2000). The spectral index of the dust echo at ν as a function of t is then given by:

$$a_\nu^E(t) = \frac{d \log F_\nu^E(t)}{d \log \nu} \quad (27)$$

$$\approx \begin{cases} a_\nu + \alpha_{\nu(1+z)} \left[1 - \frac{1}{2} \frac{t}{t_\tau} - f \tau_{\nu(1+z)} \right] & (t < t_{sc} \text{ and } t < t_\tau) \\ a_\nu + \alpha_{\nu(1+z)} \left[1 - \frac{1}{10} \frac{8-3\left(\frac{t}{t_{sc}}\right)^{-5/2}}{4-3\left(\frac{t}{t_{sc}}\right)^{-1/2}} \frac{t}{t_\tau} - f \tau_{\nu(1+z)} \right] & (t_{sc} < t < t_\tau) \\ a_\nu + \alpha_{\nu(1+z)} \left[1 - \frac{1}{10} \frac{8-3\left(\frac{t_\tau}{t_{sc}}\right)^{-5/2} \left(\frac{t}{t_\tau}\right)^{-5/4}}{4-3\left(\frac{t_\tau}{t_{sc}}\right)^{-1/2} \left(\frac{t}{t_\tau}\right)^{-1/4}} - f \tau_{\nu(1+z)} \right] & (t_{sc} < t_\tau < t) \\ a_\nu + \alpha_{\nu(1+z)} \left[\frac{1}{2} - f \tau_{\nu(1+z)} \right] & (t_\tau < t \text{ and } t_\tau \ll t_{sc}) \end{cases} \quad (28)$$

where a_ν is the spectral index of the pulse of light. This approximation holds if $f = 0$, or if $f = 1$ and $\theta_{jet} \ll \min\{\theta, (\theta/\tau)(r/R)\}$. We plot how the spectral index of the dust echo differs from the spectral index of the pulse of light as a function of time for different values of f , $R(1+z)$, and $\tau_{\nu(1+z)}$ in the bottom panels of Figures 4 – 7; however, consider the following limiting cases:

f = 0 (Figures 4 and 5, bottom panels): Since blue light scatters with a higher probability than red light, and since both the dust echo and the pulse of light are extinguished equally at

$t = 0$, the dust echo is initially bluer than the pulse of light, independent of the value of $\tau_{\nu(1+z)}$: $a_{\nu}^E(t) - a_{\nu} \approx \alpha_{\nu(1+z)} \approx 1$. If $\tau_{\nu(1+z)} \lesssim 12$, then $t_{sc} \lesssim t_{\tau}$. Consequently, the dust echo reddens to $a_{\nu}^E(t) - a_{\nu} \approx \alpha_{\nu(1+z)}(1 - \tau_{\nu(1+z)}/24)$ at $t \approx t_{sc}$. If $\tau_{\nu(1+z)} \lesssim 1.5$, then it grows redder to $a_{\nu}^E(t) - a_{\nu} \approx \alpha_{\nu(1+z)}(1 - \tau_{\nu(1+z)}/8)$ at $t \approx t_{off}$, after which it grows bluer and returns to its original color. If $1.5 \lesssim \tau_{\nu(1+z)} \lesssim 12$, then it grows redder to $0.5\alpha_{\nu(1+z)} \lesssim a_{\nu}^E(t) - a_{\nu} \lesssim 0.7\alpha_{\nu(1+z)}$ at $t \approx t_{\tau}$, after which it grows bluer to a color of $a_{\nu}^E(t) - a_{\nu} \approx 0.8\alpha_{\nu(1+z)}$ until after $t = t_{off}$. After this time, it grows bluer still and returns to its original color. If $\tau_{\nu(1+z)} \gg 12$, then t_{τ} is initially less than t_{sc} . Consequently, the dust echo reddens to $a_{\nu}^E(t) - a_{\nu} \approx 0.5\alpha_{\nu(1+z)}$ at $t \approx t_{\tau}$, after which its color does not change until after $t = t_{off}$. After this time, its color grows bluer and returns to its original color.

$\mathbf{f} = \mathbf{1}$ (Figures 6 and 7, bottom panels): The spectral index of the dust echo has the same dependence on time as in the $f = 0$ case, but is redder than the pulse of light by an additional $\tau_{\nu(1+z)}$: since the dust is destroyed along the line of sight, the pulse of light is not similarly reddened.

In any case, if $\tau_{\nu(1+z)} \lesssim 12$, then the spectral index of the dust echo does not change in time by more than $0.5\alpha_{\nu(1+z)} \approx 0.5$, and if $\tau_{\nu(1+z)} \ll 1$, then it does not change at all. If $\tau_{\nu(1+z)} \gtrsim 12$, then the dust echo is probably too extinguished to be observed; consequently, we do not explore this case in detail in this paper.

2.2. Dust Echoes from a Scattered Collimated Pulse of Light

We now model the more general case of the dust echo of a collimated pulse of light that is emitted from a distance that is much less than R . Let the pulse of light be emitted between $0 < \theta < \theta_{col}$ and $\pi - \theta_{col} < \theta < \pi$ (we assume bipolar emission), where $\theta_{col} > \theta_{jet}$. The only difference from the isotropic case is that $dP_{\nu(1+z)}^{sc}(r, \theta) = 0$ if $\theta_{col} < \theta < \pi - \theta_{col}$. Consequently, the light curve is again given by Equation (16), but with

$$P(r') = \begin{cases} 0 & [r' < r(\pi - f\theta_{jet})] \\ 1 & [r(\pi - f\theta_{jet}) < r' < r(\pi - \theta_{col})] \\ 0 & [r(\pi - \theta_{col}) < r' < r(\theta_{col})] \\ 1 & [r(\theta_{col}) < r' < r(f\theta_{jet})] \\ 0 & [r' > r(f\theta_{jet})] \end{cases}, \quad (29)$$

$$r = \begin{cases} R & [t < t(\theta_{col})] \\ r(\theta_{col}) & [t(\theta_{col}) < t < t(\pi - \theta_{col})] \\ R & [t(\pi - \theta_{col}) < t < t(\pi - f\theta_{jet})] \\ r(\pi - f\theta_{jet}) & [t > t(\pi - f\theta_{jet})] \end{cases}, \quad (30)$$

and

$$\theta = \begin{cases} \theta(R) & [t < t(\theta_{col})] \\ \theta_{col} & [t(\theta_{col}) < t < t(\pi - \theta_{col})] \\ \theta(R) & [t(\pi - \theta_{col}) < t < t(\pi - f\theta_{jet})] \\ \pi - f\theta_{jet} & [t > t(\pi - f\theta_{jet})] \end{cases} . \quad (31)$$

This interaction of the dust and light geometries is depicted in the bottom panel of Figure 3.

We plot example light curves and color histories for different values of θ_{col} and $\tau_{\nu(1+z)}$ in Figures 8 and 9, but basically, they are the same as in the isotropic case until $t \approx t(\theta_{col})$. After this time, the light curve falls off as $F_{\nu}^E(t) \propto t^{-2}$ if $t < t_{\tau}$, as $F_{\nu}^E(t) \propto t^{-7/4}$ if $t > t_{\tau}$ and $t_{sc} < t_{\tau}$ [$\tau_{\nu(1+z)} \lesssim 12r/R$], and as $F_{\nu}^E(t) \propto t^{-1}$ if $t > t_{\tau}$ and $t_{\tau} < t_{sc}$ [$\tau_{\nu(1+z)} \gg 12r/R$], because the observed light no longer includes light that scattered close to the evacuated region where the density (and the optical depth) is highest. A direct consequence of this is that the dust echo grows bluer. The light curve and color history partially recover their isotropic characteristics after $t \approx t(\pi - \theta_{col})$, when back-scattered light from the far side of the evacuated region can first reach the observer; however, the light curve falls off rapidly after $t \approx t_{off}$.

2.3. Dust Echoes from Scattered Afterglow Light

We now model the even more general case of the dust echo of collimated light that is emitted as a function of time (i.e., an afterglow) from a distance that is much less than R . Since the afterglow emits light continuously, at any time t after the burst, we receive scattered light that was emitted by the afterglow at all prior observer-frame times: $0 < t' < t$. Consequently, the dust echo light curve is given by integrating Equation (16) over these times:

$$F_{\nu}^E(t) = \frac{3c\tau_{\nu(1+z)}}{2R(1+z)} \int_0^t \int_r^{\infty} P(r') \left(\frac{r'}{R}\right)^{-3} \frac{e^{-[\tau_{\nu(1+z)}(r',\theta) - (1-f)\tau_{\nu(1+z)}]}}{(5 - 4\cos\theta)^{3/2}} d\left(\frac{r'}{R}\right) F_{\nu}(t') dt', \quad (32)$$

where $P(r')$, r , and θ are given by Equations (29), (30), and (31), respectively, but with

$$r(\theta) = \frac{c(t - t')}{(1 - \cos\theta)(1 + z)}, \quad (33)$$

$$\theta(r) = \cos^{-1} \left[1 - \frac{c(t - t')}{r(1 + z)} \right], \quad (34)$$

and

$$t(\theta) = t' + \frac{(1 - \cos\theta)R(1 + z)}{c}. \quad (35)$$

$F_{\nu}(t')$ is the light curve of the afterglow, and θ_{col} is now a function of time that we approximate by:

$$\theta_{col} = \min \left\{ (\theta_{jet}^2 + \gamma^{-2})^{1/2}, \frac{\pi}{2} \right\}. \quad (36)$$

Let t_{jet} be the time when $\gamma = \theta_{jet}^{-1}$. Then from Equation (3),

$$\gamma = \begin{cases} 10 \left(\frac{\theta_{jet}}{0.1} \right)^{-1} \left(\frac{t'}{t_{jet}} \right)^{-1/4} & (t' < t_{jet}) \\ 10 \left(\frac{\theta_{jet}}{0.1} \right)^{-1} \left(\frac{t'}{t_{jet}} \right)^{-1/2} & (t' > t_{jet}) \end{cases}, \quad (37)$$

and

$$t_{jet} = 5.5(1+z) \left[\frac{n(1 \text{ pc})}{1 \text{ cm}^3} \right]^{-1} \left(\frac{E\theta_{jet}^2}{10^{52} \text{ erg}} \right) \left(\frac{\theta_{jet}}{0.1} \right)^2 \text{ hr}. \quad (38)$$

The value of $\gamma \propto (t')^{-1/2}$ after t_{jet} (e.g., Sari, Piran, & Halpern 1999); this is true in either a $n(r) \propto r^0$ or a $n(r) \propto r^{-2}$ circumburst medium.

Finally, we model the light curve of the afterglow. In this paper, we simply take:

$$F_\nu^E(t') = \begin{cases} F_\nu^E & (t' < t_0) \\ F_\nu^E \left(\frac{t'}{t_0} \right)^{-1.2} & (t_0 < t' < t_{jet}) \\ F_\nu^E \left(\frac{t_{jet}}{t_0} \right)^{-1.2} \left(\frac{t'}{t_{jet}} \right)^{-2.4} & (t' > t_{jet}) \end{cases}, \quad (39)$$

which is typical of observed afterglows after t_0 , where t_0 may be as little as minutes or as much as hours after the burst (e.g., Sari, Piran, & Narayan 1998). Before t_0 , the light curve can depend on time in a variety of ways (e.g., Sari, Piran, & Narayan 1998); however, we do not model it any more carefully than a constant. Instead, we show that the value of t_0 has little effect on the light curve and color history of the dust echo (see Figure 13).

We plot example light curves and color histories for different values of θ_{jet} , t_{jet} , and t_0 in Figures 10 – 13. The results are similar to the case of a collimated pulse of light with collimation half-angle given by the value of θ_{col} when most of the afterglow light is emitted; i.e., by the value of θ_{jet} , since most of the afterglow light is emitted between $t_0 < t' < t_{jet}$, and $\theta_{col} \approx \theta_{jet}$ during these times.

In Figure 10, we take $f = 1$, $R(1+z) = 0.1 \text{ pc}$, $\tau_{\nu(1+z)} = 1$, $t_0 = 1 \text{ hr}$, $\theta_{jet} = 5^\circ, 10^\circ, 20^\circ$, and 40° , and $t_{jet} = 0.35, 1.4, 5.6$, and 22 dy , respectively. These values of t_{jet} are given by Equation (38) with $z = 1$, $n(1 \text{ pc}) = 1 \text{ cm}^{-3}$, $E\theta_{jet}^2 = 10^{52} \text{ erg}$, and these values of θ_{jet} . Clearly, for these values of the parameters, the dust echo of the afterglow is substantially fainter than the afterglow itself. Changing the value of $\tau_{\nu(1+z)}$ only makes the dust echo fainter relative to the afterglow: decreasing $\tau_{\nu(1+z)}$ decreases the amount of light scattered into the line of sight, and increasing $\tau_{\nu(1+z)}$ increases how much the dust echo is extinguished, but does not increase how much the afterglow is extinguished since $f = 1$. The dust echo can be made brighter than the afterglow if $f = 0$ and $\tau_{\nu(1+z)}$ is substantially increased, but in this scenario, both the afterglow and the dust echo are substantially extinguished and consequently difficult to detect. This leaves only three scenarios in which a dust echo can be brighter than an afterglow and detectable: (1) the value of $R(1+z)$ is substantially greater than 0.1 pc , in which case the dust echo peaks at a substantially later time (but with a substantially lower peak brightness), after the light curve of

the afterglow has rolled over and “gotten out of the way” of the dust echo (Figure 11); (2) the value of $(1+z)[n(1\text{ pc})]^{-1}E\theta_{jet}^2$ is substantially less than the value assumed above, in which case the light curve of the afterglow rolls over at a substantially earlier time, again “getting out of the way” of the dust echo (Figure 12); and (3) the dust echo is dominated by scattered light not from the afterglow, but rather, from the optical flash and/or optical light from the burst itself. This last scenario can be modeled simply as a collimated pulse of light (§2.2) with $\theta_{col} \gtrsim \theta_{jet}$. We show that the choice of t_0 has little effect on the light curve and color history of the dust echo of the afterglow in Figure 13.

One aspect of the dust and light geometry that we do not model in this paper is that the afterglow is not emitted from the center of the evacuated region, but rather from the head of the jet, which is moving away from the center (toward the observer) at relativistic speeds. Since $r \approx 4\gamma^2 ct/(1+z)$ (§1), the head of the jet reaches the radius r at which most of the scattering is occurring at time

$$t_r = 8.9(1+z)^2 \left(\frac{\theta_{jet}}{0.1} \right)^4 \left(\frac{t_{jet}}{1\text{ dy}} \right)^{-1} \left(\frac{r}{1\text{ pc}} \right)^2 \text{ dy}. \quad (40)$$

Our model holds as long as most of the light is emitted before some reasonable fraction of t_r . For afterglow light curves like the one described in Equation (39), most of the afterglow light is emitted before t_{jet} , by a factor of a few. Consequently, our model only holds as long as $t_r \gtrsim t_{jet}$, or

$$r \gtrsim 0.08 \left[\frac{n(1\text{ pc})}{1\text{ cm}^3} \right]^{-1} \left(\frac{E\theta_{jet}^2}{10^{52}\text{ erg}} \right) \text{ pc}. \quad (41)$$

However, if the dust echo is dominated by scattered light from the optical flash and/or optical light from the burst, as opposed to light from the afterglow, then the light is obviously emitted long before t_r , and consequently, the model holds for substantially smaller values of r as well.

3. Dust Echoes from Absorbed and Thermally Re-Emitted Light

The dust echo from absorbed and thermally re-emitted light is easier to model than the dust echo from scattered light: unlike Equation (12), the differential cross section for dust absorption has no dependence on θ :

$$\frac{d\sigma_{\nu(1+z)}^{abs}}{d\Omega} = \frac{(1 - \epsilon_{\nu(1+z)})\sigma_{\nu(1+z)}^{tot}}{4\pi}. \quad (42)$$

Consequently, the light curve of the dust echo of the afterglow is given by

$$F_{IR}^E(t) = \frac{c\tau_{\nu(1+z)}}{R(1+z)} \int_0^t \int_r^\infty P(r') \left(\frac{r'}{R} \right)^{-3} e^{-[\tau_{\nu(1+z)}(r',\theta) - (1-f)\tau_{\nu(1+z)}]} d\left(\frac{r'}{R} \right) F_{UV}(t') dt', \quad (43)$$

where $F_{IR}(t)$ is the bolometric light curve of the dust echo, and $F_{UV}(t')$ is the light curve of the afterglow, integrated over UV values of $\nu(1+z)$. If the re-radiating dust particles had a common temperature T , the spectrum of the dust echo would be thermal. However, the temperature of

the dust particles decreases with radius as $T \propto r^{-1/2}$ (Waxman & Draine 2000); consequently, the spectrum should fall off more slowly than a blackbody redward of the spectral peak. Since dust is rapidly sublimated at temperatures $T \gtrsim 2300$ K, the spectral peak should occur only at frequencies $\nu \lesssim 1.35 \times 10^{14}(1+z)^{-1}$ Hz. For a burst at $z = 0$, this is in the K band.

We plot example light curves for different values of θ_{jet} in Figure 14. However, consider the limiting cases for an isotropic pulse of light. The thermal dust echo has the same limiting cases as the scattered dust echo (e.g., Equation 23), except that $t_{sc} = \infty$. Consequently, if $t_\tau > t(\theta_{col})$, then $F_{IR}(t) \propto t^0$ until $t \approx t(\theta_{col})$, after which $F_{IR}(t) \propto t^{-2}$ until $t \approx t_\tau$. After this time, $F_{IR}(t) \propto t^{-1}$. If $t_\tau < t(\theta_{col})$, then $F_{IR}(t) \propto t^0$ until $t \approx t_\tau$, after which $F_{IR}(t) \propto t^{-1}$.

4. Can the Late-Time Afterglow of GRB 970228 Be Explained by a Dust Echo?

No. First of all, the late-time afterglow of GRB 970228 cannot be explained by a thermal dust echo, because its spectral flux distribution (Figure 1) peaks between the V and B bands in the source frame, not in or beyond the K band (§3). Secondly, the late-time afterglow of GRB 970228 cannot be explained by a dust echo due to scattered afterglow light, because its light curve (e.g., Reichart 1999) does not roll over at sufficiently early times for the dust echo to be able to outshine the afterglow when the second component to the light curve peaks, about 20 – 30 days after the burst (§2.3). This leaves only the possibility of a dust echo due to scattered light from the optical flash and/or optical light from the burst itself. To investigate this possibility, we have fitted the spectral flux distribution of Figure 1 with two power laws, one to the reddest three points and one to the bluest three points. We find that the spectral index redward of and including the source-frame V band is $a_{rV}^E = 2.29_{-0.58}^{+1.43}$, and that the spectral index blueward of and including the source-frame B band is $a_{bB}^E = -3.09_{-0.55}^{+0.54}$. The corresponding optical depths as a function of the spectral index of the pre-scattered, unextinguished light are given by Equation (28) with f set to 1. Taking $\alpha_{rV} = 1.6$ and $\alpha_{bB} = 1$ (§2.1), we find that for an intrinsic spectral index of $-1/2$, $\tau_{rV} = -0.99_{-0.93}^{+0.44}$ and $\tau_{bB} = 3.34_{-0.60}^{+0.60}$, and for an intrinsic spectral index of $1/3$, $\tau_{rV} \approx -0.47_{-0.93}^{+0.44}$ and $\tau_{bB} = 4.17_{-0.60}^{+0.60}$. Intrinsic spectral indexes of $-1/2$ and $1/3$ are typical of what one expects for the optical flash blueward and redward of the synchrotron peak, respectively, which one expects to be at optical/NIR wavelengths during the optical flash (e.g., Sari & Piran 1999). In the case of optical light from the burst itself, one expects a spectral index of $1/3$ across the optical and NIR bands (e.g., Sari & Piran 1999). Consequently, in either interpretation, the value of τ_{rV} is borderline consistent with being physical (i.e., non-negative), but the rapid transition between A_R or $A_V \approx 0$ to A_B or $A_U \approx 3.6 \pm 0.7$ or 4.5 ± 0.7 is unphysical. Consequently, we find that the late-time afterglow of GRB 970228 cannot be explained by a dust echo.

Unfortunately, the late-time afterglow of GRB 980326 was not sufficiently well sampled to distinguish between a supernova and a dust echo explanation

5. Discussion and Conclusions

We have modeled and computed dust echo light curves and spectra for dust and light geometries that are more realistic than had previously been considered. In particular, bursts are very likely to be collimated (on both theory and energetics grounds), and dust may be destroyed by the progenitor, as well as the burst, optical flash, and/or afterglow. Also, the optical flash and/or optical light from the burst itself may result in dust echoes. Consequently, dust echoes may be a way to measure the brightness of the optical flash and/or the optical brightness of the burst.

We find that dust echo spectra can be used to distinguish different types of dust echoes from each other, and from supernovae. For example, thermal dust echoes peak in the source-frame K band, while supernovae peak in the source-frame V or B bands. In the absence of extinction (e.g., in the source-frame NIR), dust echoes due to scattered light are bluer than the pre-scattered light by about 1 – 1.6 in the spectral index, and consequently, can peak in the source-frame V or B bands, or other nearby bands, depending on the amount of extinction. However, dust echoes due to scattered light do not peak as sharply as do thermal dust echoes or supernovae. It is for these reasons that we find that the late-time afterglow of GRB 970228 cannot be explained by a dust echo.

Furthermore, dust echo light curves can often be used to distinguish dust echoes from supernovae. While dust echoes can peak about $20(1+z)$ days after a burst, they also can peak substantially earlier and substantially later. However, supernovae can peak only at this time.

Finally, dust echoes can be used to probe the circumburst environment, and the effects of the progenitor, burst, optical flash, and/or afterglow on this environment. For example, if the dust echo light curve brightens, levels off for a long time, and then fades, then $f = 0$ and dust had not (yet) been destroyed along the line of sight. If the dust echo light curve brightens and fades without leveling off, then $f = 1$ and dust had been destroyed along the line of sight. The time at which the dust echo light curve begins to fade gives you (most likely) a degenerate combination of R and θ_{jet} . The color of the dust echo, particularly near peak, gives you $\tau_{\nu(1+z)}$, and consequently a degenerate combination of R and $n(1 \text{ pc})$. The time at which the afterglow light curve rolls over due to lateral expansion of the jet gives you a degenerate combination of θ_{jet} and $n(1 \text{ pc})$. Consequently, R , θ_{jet} , and $n(1 \text{ pc})$ can be separated and solved for. Furthermore, the brightness of the dust echo, particularly near peak, gives you a degenerate combination of R , $\tau_{\nu(1+z)}$, and the fluence of the pre-scattered, unextinguished light; consequently, this fluence can be solved for as well. Also, by comparing $\tau_{\nu(1+z)}$ to the same quantity measured from the afterglow, and to absorption measured in X rays, one can further probe dust destruction along the line of sight, and the metals-to-dust ratio off the line of sight where dust is not destroyed, respectively.

Support for this work was provided by NASA through the Hubble Fellowship grant #HST-SF-01133.01-A from the Space Telescope Science Institute, which is operated by the Association

of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. I am very grateful to Re'em Sari for invaluable discussions about the dust and light geometries. I am also grateful to Don Lamb for discussions about dust echoes in general.

REFERENCES

- Berger, E., et al. 2000, *ApJ*, 545, 56
- Bloom, J. S., et al. 1999, *Nature*, 401, 453
- Bloom, J. S., Djorgovski, S. G., & Kulkarni, S. R. 2001, *ApJ*, submitted (astro-ph/0008029)
- Chevalier, R. A., & Li, Z. 1999, *ApJ*, 520, L29
- Draine, B. T., & Salpeter, E. E. 1979, *ApJ*, 231, 438
- Esin, A. A., & Blandford, R. 2000, *ApJ*, 534, L151
- Galama, T. J., et al. 2000, *ApJ*, 536, 185
- Halpern, J. P., et al. 2000, *ApJ*, 543, 697
- Harrison, F. A., et al. 1999, *ApJ*, 523, L121
- Lamb, D. Q. 1997, poster at Gamma-Ray Bursts: 4th Huntsville Symposium
- Lamb, D. Q. 2000, in *Procs. of Gamma-Ray Bursts in the Afterglow Era: 2nd Workshop*, in preparation
- Lamb, D. Q., & Reichart, D. E. 2000, *ApJ*, 536, 1
- Panaiteanu, A., Mészáros, P., & Rees, M. J. 1998 *ApJ*, 503, 314
- Price, P. A., et al. 2001, *ApJ* (Letters), submitted (astro-ph/0012303)
- Reichart, D. E. 1999, *ApJ*, 521, L111
- Reichart, D. E. 2001, *ApJ*, in press (astro-ph/9912368)
- Reichart, D. E., Castander, F. J., & Lamb, D. Q. 2000, in *Cosmic Explosions: Tenth Astrophysics Conference*, AIP Conference Proceedings 522, eds. S. S. Holt and W. M. Zhang (Melville, New York: AIP), 253
- Reichart, D. E., Lamb, D. Q., & Castander, F. J. 2000, in *Gamma-Ray Bursts: 5th Huntsville Symposium*, AIP Conference Proceedings 526, eds. R. M. Kippen, R. S. Mallozzi, and G. J. Fishman (Melville, New York: AIP), 414

- Rhoads, J. E., & Fruchter, A. S. 2000, ApJ, in press (astro-ph/0004057)
- Sari, R., & Piran, T. 1999, ApJ, 520, 641
- Sari, R., Piran, T., & Halpern, R. 1999, ApJ, 519, L17
- Sari, R., Piran, T., & Narayan, R. 1998, ApJ, 497, L17
- Stanek, K. Z., et al. 1999, ApJ, 522, L39
- Waxman, E., & Draine, B. T. 2000, ApJ, 537, 796
- White, R. L. 1979, ApJ, 229, 954

Fig. 1.— The K- through V-band spectral flux distribution of the late-time afterglow of GRB 970228 after subtracting out the contribution of the host galaxy and correcting for Galactic extinction (filled circles and upper limit), and the I- through U-band spectral flux distribution of SN 1998bw after transforming to the redshift of GRB 970228, $z = 0.695$ (Bloom, Djorgovski, & Kulkarni 2001), and correcting for Galactic extinction (unfilled circles; Reichart, Castander, & Lamb 2000; Reichart, Lamb, & Castander 2000). The K-band upper limits are 2 and 3σ . We have fitted power laws to the K-, J-, and I-band observations, and the I-, R-, and V-band observations of the late-time afterglow of GRB 970228 (see §4). The solid lines are the best fits, and the dotted lines are the 1σ uncertainties in these fits. Clearly, the spectral flux distribution of the late-time afterglow of GRB 970228 is very blue redward of the peak, and very red blueward of the peak.

Fig. 2.— Dust and light geometries. Top panel: Isotropic dust destruction and an isotropic pulse of light. Dust is destroyed isotropically about the progenitor, and interior to the thick solid circle, the radius of which is $R < R_w$, where R_w is the radius of the wind termination shock. Exterior to R , $n(r) \propto r^{-2}$ out to R_w . The parabola marks the surface on which light must either scatter or be absorbed and thermally re-emitted to arrive at an observer to the far right at a constant time t after the burst, where t is given by Equation (1). Light cannot scatter or be absorbed and thermally re-emitted on the dotted portion of this surface of revolution because dust has been destroyed here. Bottom panel: Collimated dust destruction and a collimated pulse of light. Dust is destroyed within the bipolar cone (thick solid curve) out to a radius $R > R_w$. The radius R_w is marked with a thin dashed circle. Interior to R_w , $n(r) \propto r^{-2}$; exterior to R_w , $n(r) \propto r^0$. The shaded area marks the region not illuminated by the collimated pulse of light. The parabola again marks a “surface of constant time”. Light cannot scatter or be absorbed and thermally re-emitted on the dotted portions of this surface because either dust has been destroyed here, or light does not reach here. In this dust and light geometry, light clearly scatters and is absorbed and thermally re-emitted at radii $r \ll R$.

Fig. 3.— In both panels, the dust geometry is the same: dust is destroyed interior to a sphere of radius $R < R_w$ by the progenitor, and interior to a bipolar cone by the burst, optical flash, and/or afterglow (thick solid curve). Exterior to the thick solid curve, $n(r) \propto r^{-2}$ out to R_w . In the top panel, the pulse of light is isotropic; in the bottom panel, the pulse of light is collimated. The parabolas mark surfaces of constant time that are separated by factors of two in time. Clearly, these dust and light geometries would result in dust echoes with similar early-time and late-time behaviors, but different intermediate-time behaviors.

Fig. 4.— Light curves (top panel) and color histories (bottom panel) of scattering dust echoes of an isotropic pulse of light, with $f = 0$ (dust is destroyed isotropically by the progenitor, but not additionally by the burst, optical flash, and/or afterglow along the line of sight), $R(1+z) = 0.1, 0.3, 1, 3$, and 10 pc, and $\tau_{\nu(1+z)} = 0.1$. Since $f = 0$, the dust echo is bluer than the pulse of light. Since $\tau_{\nu(1+z)} \ll 1$, the color of the dust echo is relatively constant with time.

Fig. 5.— Light curves (top panel) and color histories (bottom panel) of scattering dust echoes of

an isotropic pulse of light, with $f = 0$, $R(1+z) = 0.1$ pc, and $\tau_{\nu(1+z)} = 0.1, 0.3, 1, 3, 10$. Since $f = 0$, the dust echo is bluer than the pulse of light. However, when $\tau_{\nu(1+z)} \gtrsim 1$, the dust echo does grow somewhat redder when the dust echo begins to fade away. This is due to greater extinction, due to the greater path length that the light must travel through the dust at the larger scattering angles. These larger scattering angles come into play only at late times, and are in fact responsible for the onset of the fading.

Fig. 6.— Light curves (top panel) and color histories (bottom panel) of scattering dust echoes of an isotropic pulse of light, with $f = 1$ (dust is destroyed both isotropically out to R by the progenitor, and within θ_{jet} of the line of sight out to $r > a$ few times R by the burst, optical flash, and/or afterglow), $R(1+z) = 0.1, 0.3, 1, 3$, and 10 pc, $\tau_{\nu(1+z)} = 0.1$, and $\theta_{jet} = 0.1$. The light curves “turn on” once $\theta > \theta_{jet}$. Since $\tau_{\nu(1+z)} \ll 1$, the color of the dust echo is relatively constant with time.

Fig. 7.— Light curves (top panel) and color histories (bottom panel) of scattering dust echoes of an isotropic pulse of light, with $f = 1$, $R(1+z) = 0.1$ pc, $\tau_{\nu(1+z)} = 0.1, 0.3, 1, 3, 10$, and $\theta_{jet} = 0.1$. When $\tau_{\nu(1+z)} > 1$, the dust echo is generally fainter and redder relative to the pulse of light, because the dust echo is extinguished, but the pulse of light, which travels along the dust-evacuated line of sight, is not. However, regardless of the value of $\tau_{\nu(1+z)}$, the dust echo is bright initially, when θ is only slightly greater than θ_{jet} , and again when θ is only slightly less than $\pi - \theta_{jet}$, because at these angles, scattered light travels through a minimal amount of dust before entering the dust-evacuated cone (of half-angle θ_{jet}) along the line of sight. For the same reason, the dust echo grows significantly bluer at these times.

Fig. 8.— Light curves (top panel) and color histories (bottom panel) of scattering dust echoes of a collimated pulse of light, with $f = 1$, $R(1+z) = 0.1$ pc, $\tau_{\nu(1+z)} = 0.1$, $\theta_{jet} = 0.1$ (5.7°), and $\theta_{col} = 10^\circ, 30^\circ, 50^\circ, 70^\circ$, and 90° . The $\theta_{col} = 90^\circ$ light curve and color history is the same as in Figure 7. The other light curves and color histories are the same as this one, except that the light that scattered at $\theta > \theta_{col}$ no longer contributes.

Fig. 9.— Same as Figure 8, but with $\tau_{\nu(1+z)} = 3$.

Fig. 10.— Light curves (top panel) and color histories (bottom panel) of scattering dust echoes (solid lines) of afterglow light (dotted lines), with $f = 1$, $R(1+z) = 0.1$ pc, $\tau_{\nu(1+z)} = 1$, $t_0 = 1$ hr, $\theta_{jet} = 5^\circ, 10^\circ, 20^\circ$, and 40° , and $[(1+z)/2][n(1\text{ pc})/1\text{ cm}^3]^{-1}(E\theta_{jet}^2/10^{52}\text{ erg}) = 1$. The corresponding values of t_{jet} are 0.35, 1.4, 5.6, and 22 dy, respectively (Equation 38). Clearly, the dust echoes of these afterglows would be lost in the brightness of the afterglow.

Fig. 11.— Same as Figure 10, but with $R(1+z) = 1$ pc. The dust echoes are brighter than the afterglows at times substantially later than t_{jet} .

Fig. 12.— Same as Figure 10, but with $[(1+z)/2][n(1\text{ pc})/1\text{ cm}^3]^{-1}(E\theta_{jet}^2/10^{52}\text{ erg}) = 0.1$. The corresponding values of t_{jet} are 0.035, 0.14, 0.56, and 2.2 dy, respectively. The dust echoes are

again brighter than the afterglows at times substantially later than t_{jet} .

Fig. 13.— Light curves (top panel) and color histories (bottom panel) of scattering dust echoes (solid lines) of afterglow light (dotted lines), with $f = 1$, $R(1+z) = 1$ pc, $\tau_{\nu(1+z)} = 3$, $t_0 = 0.3$, 1, and 3 hr, $\theta_{jet} = 10^\circ$, and $[(1+z)/2][n(1\text{ pc})/1\text{ cm}^3]^{-1}(E\theta_{jet}^2/10^{52}\text{ erg}) = 0.1$. The corresponding value of t_{jet} is 0.14 dy. Clearly, the choice of t_0 has little affect on the light curve and color history of the dust echo.

Fig. 14.— Light curves of thermal dust echoes (solid lines) of afterglow light (dotted lines), with $f = 1$, $R(1+z) = 1$, $\tau_{\nu(1+z)} = 3$, $t_0 = 1$ hr, $\theta_{jet} = 5^\circ$, 10° , 20° , and 40° , and $[(1+z)/2][n(1\text{ pc})/1\text{ cm}^3]^{-1}(E\theta_{jet}^2/10^{52}\text{ erg}) = 0.1$. The corresponding values of t_{jet} are 0.035, 0.14, 0.56, and 2.2 dy, respectively. Thermal dust echoes are relatively brighter at late times than scattering dust echoes, because large-angle changes in the direction of the light are just as probable as small-angle changes, but this is not the case with scattering. Large-angle changes are especially important at late times.



























